

REVISED ANALYSIS OF THE SUPERSOFT X-RAY PHASE, HELIUM ENRICHMENT, AND TURN-OFF TIME IN THE 2000 OUTBURST OF THE RECURRENT NOVA CI AQUILAE

IZUMI HACHISU

Department of Earth Science and Astronomy, College of Arts and Sciences, University of Tokyo, Komaba,
 Meguro-ku, Tokyo 153-8902, Japan
 hachisu@chianti.c.u-tokyo.ac.jp

MARIKO KATO

Department of Astronomy, Keio University, Hiyoshi, Kouhoku-ku, Yokohama 223-8521, Japan
 mariko@educ.cc.keio.ac.jp

AND

BRADLEY E. SCHAEFER

Department of Astronomy, University of Texas, Austin, TX 78712
 schaefer@astro.as.utexas.edu
to appear in the Astrophysical Journal

ABSTRACT

Recurrent nova CI Aquilae has entered the final decline phase a bit before May of 2001, about 300 days after the optical maximum, showing the slowest evolution among recurrent novae. Based on the optically thick wind mass-loss theory of the thermonuclear runaway model, we have estimated the turn-off time of the CI Aql 2000 outburst. It is in late March of 2001 after a luminous supersoft X-ray source phase lasts ~ 150 days (from November of 2000 until March of 2001). We have also obtained, by fitting our theoretical light curves with the 1917 and 2000 outbursts, the white dwarf (WD) mass to be $M_{\text{WD}} = 1.2 \pm 0.05 M_{\odot}$, the helium enrichment of the envelope is $\text{He}/\text{H} \sim 0.5$ by number, the mass of the hydrogen-rich envelope on the WD at the optical maximum is $\Delta M_{\text{max}} \sim 8.0 \times 10^{-6} M_{\odot}$, and the average mass accretion rate is $\dot{M}_{\text{acc}} \sim 1.0 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$ during the quiescent phase between the 1917 and 2000 outbursts. Using these obtained values, we have consistently reproduced the light curve in quiescence as well as of the two outbursts. We have also discussed the possibility whether or not CI Aql will explode as a Type Ia supernova in a future.

Subject headings: binaries: close — novae, cataclysmic variables — stars: individual (CI Aquilae) — stars: winds, outflows — X-rays: stars

1. INTRODUCTION AND SUMMARY

The second recorded nova outburst of CI Aquilae was discovered on 2000 April 28 UT by Takamizawa (2000), 83 years after the first recorded outburst in 1917 (Reinmuth 1925; Williams 2000). CI Aql now becomes a member of the recurrent nova class. About 300 days after the optical maximum, Hachisu & Kato (2001a) estimated various physical parameters of the CI Aql system from the light curve fitting and elucidated its nature. They derived the white dwarf (WD) mass to be $M_{\text{WD}} = 1.2 \pm 0.05 M_{\odot}$, the helium enrichment of the white dwarf envelope to be $\text{He}/\text{H} \sim 0.25$ by number, and the mass of the hydrogen-rich envelope on the white dwarf at the optical maximum to be $\Delta M_{\text{max}} \sim 5.8 \times 10^{-6} M_{\odot}$. This envelope mass indicates an average mass accretion rate of $\dot{M}_{\text{acc}} \sim 0.7 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$ during the quiescent phase between the 1917 and 2000 outbursts. They finally predicted the turn-off time of 2001 August and, therefore, that the luminous supersoft X-ray source phase lasts until August of 2001.

After Hachisu & Kato (2001a) has been published, some new observational indications appeared concerning the turn-off time of the 2000 outburst. The supersoft X-ray fluxes were too weak during June and August of 2001 (Greiner & DiStefano 2002), which indicates that the steady hydrogen shell-burning had already vanished. Schaefer (2001b) made optical photometry of CI Aql in

August of 2001 and concluded that CI Aql was $V \sim 15.3$ and almost near its quiescent phase (Mennickent & Honeycutt 1995). These results contradict the theoretical prediction that, if the hydrogen content of the white dwarf envelope is $X = 0.5$ by weight, the steady hydrogen shell-burning lasts until August of 2001 (Hachisu & Kato 2001a). Schaefer (2001c) found on 12 Harvard College Observatory archival photographs the 1941 outburst, the brightness of which is $B \sim 13.8$ on May 2 of 1941, and suggested that the 1917 discovery could indicate a recurrence timescale of about 20 yrs, with missed outbursts around 1960 and 1980. In this paper, therefore, we reanalyze the light curves of CI Aql, especially reproducing the orbital light curves newly obtained in April/May and August of 2001 (see also Schaefer 2001a,b).

Additionally, Honeycutt (2002, private communication) has revised Mennickent & Honeycutt's (1995) photometric data of CI Aql in quiescence. The revised data show about 0.62 mag down in the V -light curve so that the quiescent level is as low as $V \sim 16.1$, 0.8 mag fainter than August of 2001. This value is well consistent with Szkody's (1994) $V \sim 16.2$ in quiescence (see also Table 2 of Mennickent & Honeycutt 1995). In this sense, CI Aql did not reach its quiescent level but still showed an activity even in August of 2001. Therefore, taking these new things into account, we have fully reconsidered the CI Aql model.

To summarize the main revised results,

1. The estimated white dwarf mass is the same as the previous result, i.e., $M_{\text{WD}} = 1.2 \pm 0.05 M_{\odot}$.
2. Helium enrichment of the white dwarf envelope is estimated to be $\text{He}/\text{H} \sim 0.5$ by number, much larger than the previous value.
3. Optically thick winds stopped in early November of 2000 and the estimated turn-off time is not in August of 2001 but earlier than May of 2001, i.e., March or April of 2001. The luminous supersoft X-ray source phase lasted from November of 2000 until March or April of 2001.
4. The mass of the hydrogen-rich envelope on the white dwarf at the optical maximum is more massive, i.e., $\Delta M_{\text{max}} \sim 8.0 \times 10^{-6} M_{\odot}$ than the previous result. This envelope mass indicates an average mass accretion rate of $\dot{M}_{\text{acc}} \sim 1.0 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$ between the 1917 and 2000 outbursts.
5. These satisfy the conditions of Type Ia supernova explosion if the white dwarf consists of carbon and oxygen.

In §2, we describe our basic model of CI Aql in the late phase of the 2000 outburst and in quiescence. In §3, we have theoretically modeled orbital light curves of CI Aql in the late phase of the 2000 outburst. In §4, the 1917 outburst is numerically re-analyzed, and we re-determine the mass of the white dwarf. In §5, the 2000 outburst is also numerically re-analyzed and the duration of the strong, optically thick wind and the supersoft X-ray source phase are again estimated. Discussion follows in §6.

2. BASIC MODEL IN LATE PHASE

Our binary model is illustrated in Figure 1, which consists of a main-sequence star (MS) filling its Roche lobe, a white dwarf (WD) photosphere, and an accretion disk. A circular orbit is assumed. We also assume that the surfaces of the white dwarf, the main-sequence companion, and the accretion disk emit photons as a blackbody at a local temperature which varies with position. For the basic structure of the accretion disk, we assume an axisymmetric structure with the size and thickness of

$$R_{\text{disk}} = \alpha R_1^*, \quad (1)$$

and

$$h = \beta R_{\text{disk}} \left(\frac{\varpi}{R_{\text{disk}}} \right)^{\nu}, \quad (2)$$

where R_{disk} is the outer edge of the accretion disk, R_1^* is the effective radius of the inner critical Roche lobe for the white dwarf component, h is the height of the surface from the equatorial plane, and ϖ is the distance from the equatorial plane from the center of the white dwarf. Here, we adopt ϖ -square law ($\nu = 2$) to mimic the effect of flaring-up at the rim of the accretion disk (e.g., Schandl, Meyer-Hofmeister, & Meyer 1997). These two parameters of α and β are determined by light curve fittings.

As the August 2001 orbital light curves show non axisymmetric shape, we introduce an asymmetric configuration of the accretion disk as done by Schandl et al. (1997). They explained an asymmetric feature in the orbital light curves of the LMC luminous supersoft X-ray source CAL 87. The cause is due to the gas stream from the companion hitting the edge of the accretion disk to make a vertical spray when the mass transfer rate is as high as in the luminous supersoft X-ray sources. We have also introduced the same type of asymmetry of the accretion disk

as Schandl et al.'s, i.e.,

$$s = \frac{1 - \zeta_{\text{low}}}{2\pi - \phi_2 + \phi_1}, \quad (3)$$

$$\zeta_{\text{edge}} = \begin{cases} 1 - s(\phi - \phi_2), & \text{for } \phi_2 \leq \phi < 2\pi \\ \zeta_{\text{low}} + (1 - \zeta_{\text{low}}) \frac{\phi - \phi_1}{\phi_2 - \phi_1}, & \text{for } \phi_1 \leq \phi < \phi_2 \\ 1 - s(2\pi - \phi_2 + \phi), & \text{for } 0 \leq \phi < \phi_1 \end{cases} \quad (4)$$

and

$$\frac{z}{h} = \begin{cases} \zeta_{\text{low}} + (\zeta_{\text{edge}} - \zeta_{\text{low}}) \frac{\xi - f_a}{1 - f_a}, & \text{for } \xi \geq f_a, \\ \zeta_{\text{low}}, & \text{for } \xi < f_a, \end{cases} \quad (5)$$

where

$$\xi = \frac{\varpi}{R_{\text{disk}}}, \quad (6)$$

and ζ_{low} is a parameter specifying the degree of asymmetry, ϕ_1 is the starting angle of the vertical spray, ϕ_2 is the angle where the vertical spray reaches its maximum height, f_a is the starting radius ratio from where the vertical spray is prominent, z is the height of the disk surface from the equatorial plane for the asymmetric case. Here, we assume that the accretion disk is symmetric inside $\varpi < 0.8 R_{\text{disk}}$ ($f_a = 0.8$), and $\zeta_{\text{low}} = 0.25$, $\phi_1 = (17/16)\pi$, $\phi_2 = (9/8)\pi$, unless otherwise specified. Such an example of the accretion disk is shown in Figure 1.

The accretion luminosity of the white dwarf (e.g., Starfield, Sparks, & Shaviv 1988),

$$L_{\text{WD}} = L_{\text{WD},0} + \frac{1}{2} \frac{GM_{\text{WD}} \dot{M}_{\text{acc}}}{R_{\text{WD}}}, \quad (7)$$

is introduced, where L_{WD} is the total luminosity of the white dwarf, $L_{\text{WD},0}$ is the intrinsic luminosity of the white dwarf, and R_{WD} is the radius of the white dwarf (e.g., $R_{\text{WD}} = 0.0068 R_{\odot}$ for $1.2 M_{\odot}$ WD). The accretion luminosity is as large as $300 L_{\odot}$ for a mass accretion rate of $\dot{M}_{\text{acc}} \sim 1 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$ onto a $1.2 M_{\odot}$ WD, although it is negligibly smaller than the luminosity of steady hydrogen shell-burning during the outburst, $L_{\text{WD}} \sim 26,000 - 54,000 L_{\odot}$, for a $1.2 M_{\odot}$ WD.

The viscous luminosity of the accretion disk is also introduced. Then, the original disk surface temperature is given by

$$\sigma T_{\text{disk,org}}^4 = \frac{3GM_{\text{WD}} \dot{M}_{\text{acc}}}{8\pi \varpi^3}. \quad (8)$$

The temperature of the accretion disk surface is as high as $T_{\text{disk,org}} \sim 8500 \text{ K}$ at the radius $\varpi = 1 R_{\odot}$ for $M_{\text{WD}} = 1.2 M_{\odot}$ and $\dot{M}_{\text{acc}} = 1 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$.

The accretion disk is heated by radiation from a luminous white dwarf. Such an irradiation effect is calculated as

$$\sigma T_{\text{disk}}^4 \approx \eta_{\text{DK}} \frac{L_{\text{WD}}}{4\pi \varpi^2} \cos \theta + \frac{3GM_{\text{WD}} \dot{M}_{\text{acc}}}{8\pi \varpi^3}, \quad (9)$$

where η_{DK} is the efficiency of the irradiation and $\cos \theta$ is the inclination angle of the surface against light ray from the white dwarf (e.g., Schandl et al. 1997). We assume $\eta_{\text{DK}} = 0.5$, unless otherwise specified. This means that 50% of the absorbed radiation energy is reemitted from the disk surface while the residual 50% is converted to thermal energy that eventually heats the white dwarf. The irradiation luminosity (or temperature) is much higher than

the accretion luminosity (or temperature) during the hydrogen shell burning, i.e., $T_{\text{disk}} \sim 35,000$ K at the radius $\varpi = 1 R_{\odot}$ for $M_{\text{WD}} = 1.2 M_{\odot}$, $\cos \theta = 0.1$, and $L_{\text{WD}} = 26,000 L_{\odot}$.

The outer rim of the accretion disk is not irradiated by the white dwarf photosphere so that the temperature of the disk rim is simply assumed to be uniform, $T_{\text{disk,rim}}$, which is one of the parameter determined by fitting.

The surface temperature of the irradiated main-sequence (MS) companion is roughly estimated by

$$\sigma T_{\text{MS}}^4 \approx \eta_{\text{MS}} \frac{L_{\text{WD}}}{4\pi r^2} \cos \theta + \sigma T_{\text{MS,org}}^4, \quad (10)$$

where η_{MS} is the efficiency of the irradiation, r is the distance from the white dwarf, and $\cos \theta$ is the inclination angle of the surface. The irradiated temperature is as high as $T_{\text{MS}} \sim 26,000$ K for $\eta_{\text{MS}} = 0.5$, $\cos \theta = 0.5$, $M_{\text{WD}} = 1.2 M_{\odot}$ ($L_{\text{WD}} = 26,000 L_{\odot}$), and $r \sim 4R_{\odot}$. Therefore, the effect of the irradiation becomes very important when the irradiated hemisphere faces toward the Earth.

The temperature of each surface patch is determined by calculating all contributions from every patch of the white dwarf surface (see Hachisu & Kato 2001b, for more detail). We have divided the white dwarf surface into 16 pieces ($\Delta\theta = \pi/16$) in the latitudinal angle and into 32 pieces ($\Delta\phi = 2\pi/32$) in the longitudinal angle while the surface of the main-sequence companion is split into 32×64 pieces, respectively. The surface of the accretion disk is divided into 32 pieces logarithmically evenly in the radial direction and into 64 pieces evenly in the azimuthal angle. The outer edge of the accretion disk is also divided into 64 pieces in the azimuthal direction and 4 pieces in the vertical direction by rectangles. When the photosphere of the white dwarf becomes very small, e.g., $R_{\text{disk}}/R_{\text{ph,WD}} > 10$, we attribute the first 16 meshes to the outer region (from $\varpi = R_{\text{disk}}$ to $\varpi = R_{\text{disk}}/\sqrt{10}$) to avoid coarse meshes in the outer part, and then 16 meshes to the inner region (from $\varpi = R_{\text{disk}}/\sqrt{10}$ to $\varpi = R_{\text{ph}}$), each region of which is divided logarithmically evenly. The efficiencies of irradiation of the main-sequence companion and of the accretion disk are assumed to be $\eta_{\text{MS}} = 0.5$ and $\eta_{\text{DK}} = 0.5$, respectively.

3. LATE PHASE ORBITAL LIGHT CURVES

To detect the end of the steady hydrogen shell-burning, we have used photometry of CI Aql in May and August of 2001 (see Figs. 2-4). All photometry was with the 2.1-m telescope at McDonald Observatory with the IGI detector through a V-filter. The detector is a standard Tektronix 1026 $\times$ 1024 CCD chip while the filter is the usual Schott glass sandwich as prescribed by Bessell (1990) hence resulting in a close match to the Johnson V-filter. Standard IRAF procedures and programs were used to get differential photometry of CI Aql with respect to several calibration stars from Henden & Honeycutt (1995). A total of 118 and 429 magnitudes were obtained in April/May and August 2001, with a typical photometric uncertainty of 0.02 mag. The photometric data are wrapped in one orbital phase, with a new ephemeris obtained by Matsumoto et al. (2001), i.e.,

$$t(\text{HJD}) = 2,451,701.2086 + 0.61835 \times E, \quad (11)$$

at eclipse minima.

The orbital light curve of CI Aql in quiescence were obtained by Mennickent & Honeycutt (1995). Recently, Honeycutt (2002, private communication) rectified the data. Essential points are that their new and old orbital light curves are nearly indistinguishable in shape but the new data are about 0.6 mag fainter than the old ones. We have shown these two light curves in Figures 2-4.

To fit our theoretical light curves with the orbital photometry, we have calculated V light curves by changing the parameters of $\alpha = 0.5$ –2.0 by 0.1 step, $\beta = 0.05$ –0.50 by 0.05 step, $T_{\text{MS,org}} = 5000$ –10000 K by 100 K step, $T_{\text{disk,rim}} = 4000$ –8000 K by 100 K step, $L_{\text{WD},0} = 0$ –26,000 $L_{\odot}$ by 100 $L_{\odot}$ step, and $i = 70$ –90° by 1° step and seek the best-fit model by the visual comparison of the model with the data. After obtaining the best fit model for symmetric configurations, we further seek the best fit model for August 2000 data by changing the asymmetric configuration parameter, ζ_{low} , from 1.0 to 0.05 by 0.05 step. The best fit asymmetric models are also shown in Figures 2-4 only for August 2000 data. Dashed lines indicate the symmetric case while solid lines denote the asymmetric case. The adopted parameters are tabulated in Table 1. Enlargement around the eclipse is also shown in Figure 5 only for the $1.5M_{\odot}$ MS companion.

We have calculated the absolute M_B, M_V , and M_I magnitudes of our model (see Hachisu & Kato 2001b, in more detail). The direct fit with the observational apparent magnitude, m_V , gives us a distance modulus, $(m - M)_V$, as shown in Figures 2-4. Since each direct fit determine a distance modulus, we have several different distance moduli for each light curve. We have to choose the most appropriate distance modulus among them. This value is iteratively sought in such a way that one distance modulus is consistent all among the light curves of the late phase (May/April of 2001), the quiescent phase, the 1917 outburst, and the 2000 outburst as well as the color excess of $E(B - V)$, all of which will be discussed later in more detail.

The light curves are calculated for three companion masses of $M_{\text{MS}} = 1.0 M_{\odot}$, $1.5 M_{\odot}$ and $2.0 M_{\odot}$ as shown in Figures 2-4. Since we have obtained similar light curves for these three cases, we cannot determine the companion mass without additional information. One may distinguish them from the original temperature of the companion in quiescence; it is too high in the case of $1.0 M_{\odot}$ (7,500–7,700 K vs. $T_{\text{ZAMS}} = 5,600$ K) and too low in the case of $2.0 M_{\odot}$ (6,900–6,700 K vs. $T_{\text{ZAMS}} = 9,100$ K) compared with their zero-age main-sequence (ZAMS) surface temperature (Bressan et al. 1993). Therefore, we adopt $M_{\text{MS}} = 1.5 M_{\odot}$ in the following discussion.

The mass transfer is as high as $\sim 1 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$, and this indicates a thermally unstable mass transfer (van den Heuvel et al. 1992), which requires $M_{\text{MS}}/M_{\text{WD}} \gtrsim 1.1$ (Webbink 1985). Moreover, of our best-fit model, the original surface temperature of the main-sequence companion is roughly consistent with the effective temperature of a $\sim 1.5 M_{\odot}$ zero-age main-sequence star ($T_{\text{MS,org}} = 7,300$ –7,100 K vs. $T_{\text{ZAMS}} = 7200$ K). Since the radius of the $1.5 M_{\odot}$ ZAMS star is about $1.45 R_{\odot}$, this $1.5 M_{\odot}$ main-sequence companion has expanded to fill the Roche lobe of $\sim 1.7 R_{\odot}$ after the central hydrogen has decreased

from $X = 0.7$ at the zero-age to $X = 0.5$ at the age of ~ 1 G yr (Bressan et al. 1993). If the companion has already lost significant mass to the white dwarf component, its original zero-age mass might be more massive than $1.5 M_{\odot}$.

The edge of the accretion disk is not fully occulted by the companion star at eclipse minima as shown in Figure 6. This partial occultation implies a shift of the phase position at eclipse minima. It may advance when the asymmetry of the accretion disk is relatively large and the inner side of the spray is brightly illuminated by the white dwarf. However, our best fit model in Figure 5 shows no significant advance in the orbital phase at eclipse minima mainly because the asymmetry of $\zeta_{\text{low}} = 0.7$ is not so large.

We are able to estimate the turn-off time of steady hydrogen shell-burning on the white dwarf. The orbital light curve fitting for May 2001 suggests us that the intrinsic luminosity of the white dwarf has already decreased to about $L_{\text{WD},0} \sim 11,000 L_{\odot}$ from $26,000 L_{\odot}$ for the lower limit of steady hydrogen shell-burning. The orbital light curve fitting for August of 2001 indicates that the intrinsic luminosity of the white dwarf has much decreased to about $L_{\text{WD},0} \sim 2,200 L_{\odot}$. Thus, we have estimated the turn-off time a bit before May of 2001, that is, March or April of 2001. After the turn-off time, the white dwarf began to cool down. It should be noted, however, that hydrogen shell burning does not vanish suddenly, but gradually extinguishes. Then, our theoretical definition of the turn-off time is when the nuclear luminosity decreases below the diffusive luminosity of the white dwarf. In other words, the diffusive luminosity of the white dwarf is almost balanced with the nuclear luminosity before the turn-off time. Nuclear burning still partly contributes to the luminosity of the white dwarf even after the turn-off time. A large part of the luminosity in the cooling phase comes from the thermal energy in the hydrogen-rich envelope and the helium shell (ash of hydrogen burning) of the white dwarf.

We have calculated the color index to estimate the color excess of CI Aql. A rather blue color index of $(B - V)_c \approx 0.0$ is obtained for the case of a $1.5 M_{\odot}$ companion as shown in Figure 7. This suggests a large color excess of $E(B - V) = (B - V)_o - (B - V)_c \approx 1.0$ with the observed color of $(B - V)_o \sim 1.0$ (see Table 2 of Mennickent & Honeycutt 1995), where $(B - V)_c$ is the calculated color and $(B - V)_o$ is the observed color. This color excess is roughly consistent with $E(B - V) = 0.85 \pm 0.3$ estimated by Kiss et al. (2001). Therefore, we adopt $E(B - V) = 1.0$ in this paper.

4. THE 1917 OUTBURST

The light curve of the 1917 outburst has also been re-analyzed with the newly obtained system parameters. The Tycho B magnitude of the 1917 outburst has been reported by Williams (2000), which is shown in Figure 8 together with our calculated B light curves. We have obtained the apparent distance modulus of $(m - M)_B = 15.06$ based on the color excess of $E(B - V) = 1.0$ and the V -magnitude distance modulus estimation of $(m - M)_V = 14.06$. The numerical method for obtaining light curves is the same as that in our previous one (Hachisu & Kato 2001a) and it is fully described in Hachisu & Kato (2001b).

Assuming a binary system consisting of a white dwarf

with the mass of $M_{\text{WD}} = 1.0, 1.1, 1.15, 1.2, 1.3$, and $1.377 M_{\odot}$ and a $1.5 M_{\odot}$ lobe-filling, main-sequence companion, we have calculated the developing of the outburst and obtained B light curves for the hydrogen content of $X = 0.7$. In Figure 8, we do not include the effect of the accretion disk or the irradiation effect of the companion. These irradiation effects do not contribute much to the B light until the B -magnitude decreases to $B \sim 12$ mag, since the white dwarf photosphere has expanded larger than the binary orbit (Kato & Hachisu 1994). It is clear that there are no mid plateau phases around $V \sim 13.5$ for the case of no accretion disk and no irradiation of the companion.

The white dwarf mass is again determined to be $M_{\text{WD}} = 1.2 \pm 0.05 M_{\odot}$ from the fitting as seen in Figure 8. The various physical parameters of the outbursts are summarized in Table 2. Here, Z is the content of heavy elements by weight, ΔM_{max} is the hydrogen-rich envelope mass at the optical maximum, η_{wind} is the ratio of the mass lost by winds, ΔM_{wind} , to ΔM_{max} , i.e., $\eta_{\text{wind}} = \Delta M_{\text{wind}} / \Delta M_{\text{max}}$, t_2 is the time taken to drop 2 mag from maximum, t_3 is the time to drop 3 mag from maximum, t_{wind} is the duration of optically thick wind phase, and t_{H} is the duration of steady hydrogen shell-burning.

We are not able to determine the hydrogen content only from the fitting with the 1917 light curve, because much lower contents of hydrogen such as $X = 0.5$ and $X = 0.35$ by weight give a similar decline rate (t_2 or t_3) of the early phase light curve as listed in Table 2.

5. THE 2000 OUTBURST

CI Aql erupted in 2000 April and was densely observed in various optical bands (e.g., Kiss et al. 2001; Matsumoto et al. 2001). We have determined the hydrogen content of the white dwarf envelope from the turn-off time of the 2000 outburst.

We summarize the global feature of the optical light curve: its optical maximum ($V \sim 9$ mag) was reached on 2000 May 5 (HJD 2451669.5, Kiss et al. 2001). As shown in Figure 9, the visual brightness quickly decreased to 13.5 mag in about 60 days. Then, it stays at $V \sim 14$ mag, i.e., a plateau phase. This mid plateau phase is very similar to that of U Sco and can also be explained by the irradiation of the accretion disk (Hachisu et al. 2000a,b; Thoroughgood et al. 2001). Therefore, we have reproduced the light curve in the plateau phase by assuming model parameters similar to those of U Sco.

Theoretical light curves in Figure 9 are calculated for a pair of a $1.2 M_{\odot}$ white dwarf and a $1.5 M_{\odot}$ main-sequence companion. In the plateau phase, the light curve is determined mainly by the irradiations of the accretion disk and the main-sequence companion, because the white dwarf photosphere becomes much smaller than the binary size. The unheated surface temperatures of the companion are fixed to be $T_{\text{MS,org}} = 7,300$ K throughout the 2000 outburst and $T_{\text{disk,rim}} = 6,600$ K at the disk rim, which are the same as the model parameters in the late phase of the 2000 outburst.

The luminosity of the accretion disk depends on both the thickness β and the size α . Here, we change α by 0.1 step and β by 0.05 step and seek the best fit model for the wind phase, the steady hydrogen shell-burning phase,

and the cooling phase. Finally we have adopted a set of $\alpha = 3.0$ and $\beta = 0.05$ during the wind phase to reproduce the orbital modulation observed by Matsumoto et al. (2001), $\alpha = 0.8$ and $\beta = 0.30$ in the steady hydrogen shell-burning phase, and $\alpha = 0.8$ and $\beta = 0.20$ in the cooling phase. Here, we adopt symmetric configurations of the accretion disk throughout the 2000 outburst.

The decline of the early phase ($t \sim 0$ –70 days) hardly depends on the hydrogen content, X , of the white dwarf envelope, but the ends of the wind phase and of the hydrogen shell-burning phase depend sensitively on the hydrogen content as tabulated in Table 2. Therefore, we have finally determined the hydrogen content of $X = 0.35$ and shown calculated light curves in Figure 9 for three different bands of B , V , and I_c .

Here, we have used the apparent distance modulus of $(m - M)_V = 14.06$ for the V band. Adopting the color excess of $E(B - V) = 1.0$, we obtain the apparent distance moduli of $(m - M)_B = 15.06$ and $(m - M)_I = 12.46$ for B and I_c bands, respectively. Here, we adopted the relation of $E(I - V)/E(B - V) = -1.60$ (e.g., Rieke & Lebofsky 1985). Both the B and I_c light curves are in good agreement with the observational points as shown in Figure 9. This indicates the consistency of our estimated values. Various physical values of the 2000 outburst are summarized in Table 3.

Matsumoto et al. (2001) reported a sharp ~ 1 mag drop of R_c -magnitude on 23 November 2000 and Kiyota (2001, VSNET archives, <http://vsnet.kusastro.kyoto-u.ac.jp/vsnet/>) also observed a ~ 1.5 mag drop of I_c -magnitude around the same day as shown in Figure 9. If we attribute this drop to the end of wind phase, the hydrogen content of $X = 0.35$ is consistent with the drop.

We have re-estimated the luminous supersoft X-ray source phase. No luminous supersoft X-rays are expected in the early decline phase because the photospheric radius of the white dwarf is rather large and the photospheric temperature is relatively low. After the recurrent nova enters the plateau phase, the photospheric temperature of the white dwarf increases enough to emit supersoft X-rays (Kato 1999). During the massive wind phase, however, we do not expect supersoft X-rays because they are self-absorbed by the wind itself. U Sco was observed as a luminous supersoft X-ray source in the mid plateau phase of the 1999 outburst just after the massive wind stopped (Kahabka et al. 1999; Hachisu et al. 2000a). Our analysis suggests that the massive wind stopped before 23 November 2000. This means that a luminous supersoft X-ray phase started from November of 2000 and continued until March of 2001 as shown in Figure 9. The rather low X-ray fluxes in June and August of 2001 (Greiner & DiStefano 2002) are also very consistent with our analysis.

6. DISCUSSION

The apparent distance modulus is obtained to be $(m - M)_V = 14.06$ by fitting for the late phase of the 2000 outburst. The color excess is also estimated to be $E(B - V) = 1.0$ from the difference between the observed color and the calculated color, and the absorption of $A_V = 3.1$ from a relation of $A_V = 3.1 E(B - V)$ (e.g., Rieke & Lebofsky 1985). Therefore, the distance to CI Aql is derived to be 1.55 kpc. The distance modulus of $(m - M)_B = 15.06$ is derived from the distance modulus of $(m - M)_V = 14.06$

and the color excess of $E(B - V) = 1.0$, which are also consistent with the B light curve of the 1917 outburst. This set of $(m - M)_V = 14.06$ and $E(B - V) = 1.0$ are also in very good agreement with the V , B , and I_c light curves of the 2000 outburst as shown in Figure 9.

The orbital light curve in August of 2001 shows a significant asymmetry. This indicates that the shape of the accretion disk is changing in time at least in the final decay phase of the 2000 outburst. Assuming the asymmetry of $\zeta_{\text{low}} = 0.7$ or 0.75 , we have calculated a V light curve and added it to Figures 2-4 (solid line for August of 2001). On the other hand, the disk shape is almost symmetric for May of 2001 and in quiescence. It should be noted that the model light curve of August 2001 does not reproduce the observational lack of any secondary eclipses even if we introduce asymmetry of the accretion disk shape. It may stem from large time variations of the edge of the accretion disk.

The envelope mass at the visual maximum is estimated to be $\Delta M_{\text{max}} = 8.0 \times 10^{-6} M_\odot$ for the hydrogen content of $X = 0.35$, indicating the average mass transfer rate of $\dot{M}_{\text{acc}} = 1.0 \times 10^{-7} M_\odot \text{ yr}^{-1}$ between the 1917 and 2000 outbursts, or $\dot{M}_{\text{acc}} = 1.3 \times 10^{-7} M_\odot \text{ yr}^{-1}$ between the 1941 and 2000 outbursts. This mass accretion rate is a bit higher but almost consistent with our CI Aql model in quiescence (see Figs. 2-4).

On the other hand, if the mean recurrence period is 20 yrs as suggested by Schaefer (2001c), the average mass transfer rate increases to $\dot{M}_{\text{acc}} \sim 4 \times 10^{-7} M_\odot \text{ yr}^{-1}$. This value is high enough to maintain steady hydrogen shell-burning for hydrogen content of $X = 0.7$ on a $1.2 M_\odot$ WD when the mass transfer is steady. In other words, it is too high to be compatible with our thermonuclear runaway model, unless the mass transfer rate itself rapidly increased just before the outburst. When the hydrogen content of the transferred matter is as low as $X \sim 0.35$, the lower limit to steady hydrogen shell-burning becomes $8 \times 10^{-7} M_\odot \text{ yr}^{-1}$ (see e.g. eq.(7) in Hachisu & Kato 2001b). Then, the mass accretion rate of $\dot{M}_{\text{acc}} \sim 4 \times 10^{-7} M_\odot \text{ yr}^{-1}$ can still cause shell-flashes on a $1.2 M_\odot$ WD, although the orbital modulation of the light curve in quiescence is not consistent with such a high mass transfer rate as shown in Figures 2-4.

About 82% of the envelope mass is lost by the wind ($\Delta M_{\text{wind}} = 6.6 \times 10^{-6} M_\odot$), while the residual 18% ($\Delta M_{\text{He}} = 1.4 \times 10^{-6} M_\odot$) is left and added to the helium layer. The net mass increasing rate of the white dwarf is about $\dot{M}_{\text{He}} = 1.7 \times 10^{-8} M_\odot \text{ yr}^{-1}$ between the 1917 and 2000 outburst or $\dot{M}_{\text{He}} = 2.4 \times 10^{-8} M_\odot \text{ yr}^{-1}$ between the 1941 and 2000 outburst. These satisfy the conditions of Type Ia supernova explosion if the white dwarf consists of carbon and oxygen (Nomoto & Kondo 1991). Evolutionary paths to Type Ia supernovae via recurrent novae have been discussed in more detail in Hachisu et al. (1999a,b).

We thank the VSNET members who observed CI Aql and R. K. Honeycutt for his providing us the unpublished data of CI Aql in quiescence. This research has been supported in part by the Grant-in-Aid for Scientific Research (11640226) of the Japan Society for the Promotion of Science.

REFERENCES

- Bessell, M. S. 1990, *PASP*, 102, 1181
 Bressan, A., Fagotto, F., Bertelli, G., & Chiosi, C. 1993, *A&AS*, 100, 647
 Greiner, J., & DiStefano, R. 2002, *ApJ*, in press (astro-ph/0210035)
 Hachisu, I., & Kato, M. 2001a, *ApJ*, 553, L161
 Hachisu, I., & Kato, M. 2001b, *ApJ*, 558, 323
 Hachisu, I., Kato, M., Kato, T., & Matsumoto, K. 2000a, *ApJ*, 528, L97
 Hachisu, I., Kato, M., Kato, T., Matsumoto, K., & Nomoto, K. 2000b, *ApJ*, 534, L189
 Hachisu, I., Kato, M., & Nomoto, K. 1999a, *ApJ*, 522, 487
 Hachisu, I., Kato, M., Nomoto, K., & Umeda, H. 1999b, *ApJ*, 519, 314
 Henden, A. A., & Honeycutt, R. K. 1995, *PASP*, 107, 324
 Kahabka, P., Hartmann, H. W., Parmar, A. N., & Negueruela, I. 1999, *A&A*, 374, L43
 Kato, M. 1999, *PASJ*, 51, 525
 Kato, M., & Hachisu, I., 1994, *ApJ*, 437, 802
 Kiss, L. L., Thomson, J. R., Ogloza, W., Fűrész, G., & Sziládi, K. 2001, *A&A*, 366, 858
 Matsumoto, K., et al. 2001, *A&A*, 378, 487
 Mennickent, R. E., & Honeycutt, R. K. 1995, *Inf. Bull. Variable Stars*, 4232
 Nomoto, K., & Kondo, Y. 1991, *ApJ*, 367, L19
 Rieke, G. H., & Lebofsky, M. J. 1985, *ApJ*, 288, 618
 Reinmuth, K. 1925, *AN*, 225, 385
 Schaefer, B. E. 2001a, *IAU Circ.* 7621
 Schaefer, B. E. 2001b, *IAU Circ.* 7687
 Schaefer, B. E. 2001c, *IAU Circ.* 7750
 Schandl, S., Meyer-Hofmeister, E., & Meyer, F. 1997, *A&A*, 318, 73
 Starrfield, S., Sparks, W. M., & Shaviv, G. 1988, *ApJ*, 325, L35
 Szkody, P. 1994, *AJ*, 108, 639
 Takamizawa, K. 2000, *IAU Circ.*, 7409
 Thoroughgood, T. D., Dhillon, V. S., Littlefair, S. P., Marsh, T. R., & Smith, D. A. 2001, *MNRAS*, 327, 1323
 van den Heuvel, E. P. J., Bhattacharya, D., Nomoto, K., & Rappaport, S. 1992, *A&A*, 262, 97
 Webbink, R. F. 1985, *Interacting binary stars*, eds. J. E. Pringle & R. A. Wade (Cambridge: Cambridge Univ. Press), p.39
 Williams, D. B. 2000, *Inf. Bull. Variable Stars*, 4904

TABLE 1
 FITTED SYSTEM PARAMETERS OF CI AQUILAE^a

parameter	symbol	$M_2 = 1.0 M_\odot$	$M_2 = 1.5 M_\odot$	$M_2 = 2.0 M_\odot$
inclination angle	i	76°	74°	72°
WD luminosity (May of 2001)	$L_{WD,0}$	$16,000 L_\odot$	$11,000 L_\odot$	$7,000 L_\odot$
WD luminosity (August of 2001)	$L_{WD,0}$	$3,500 L_\odot$	$2,200 L_\odot$	$1,600 L_\odot$
MS temperature (May/August of 2001)	$T_{MS,org}$	7,500 K	7,300 K	6,900 K
MS temperature (quiescence)	$T_{MS,org}$	7,700 K	7,100 K	6,700 K
color excess	$E(B - V)$	...	1.0	...
distance modulus	$(m - M)_V$	...	14.06	...
absorption	A_V	...	3.1	...
distance	d	...	1.55 kpc	...

^aThe white dwarf mass is assumed to be $M_{WD} = 1.2 M_\odot$

TABLE 2
THEORETICAL DECLINE RATES OF LIGHT CURVES FOR THE CI AQL 1917 OUTBURST

M_{WD} ($M_{\odot}$)	X^{a}	$\eta_{\text{wind}}^{\text{b}}$	t_2 (day)	t_3 (day)	t_{wind} (day)	t_{H} (day)
1.377	0.7	0.95	3.5	5	49	59
1.3	0.7	0.90	9	14	148	287
1.2	0.35	0.82	20	35	186	310
1.2	0.5	0.83	21	37	219	450
1.2	0.7	0.84	22	39	266	720
1.15	0.7	0.82	28	48	330	1020
1.1	0.7	0.82	34	61	429	1450
1.0	0.7	0.82	55	78	730	1540

^aSolar metallicity is assumed, i.e., $Z = 0.02$.

^bThe ratio of the mass lost by winds to the accretion mass, i.e., $\Delta M_{\text{wind}}/\Delta M_{\text{max}}$.

TABLE 3
THEORETICAL PROPERTIES OF THE CI AQL 2000 OUTBURST

M_{WD} ($M_{\odot}$)	X	Z^{a}	ΔM_{max} ($M_{\odot}$)	ΔM_{wind} ($M_{\odot}$)	η_{wind}	ΔM_{He} ($M_{\odot}$)	t_2 (day)	t_3 (day)	t_{wind} (day)	t_{H} (day)
1.2	0.35	0.02	8.0×10^{-6}	6.6×10^{-6}	0.82	1.4×10^{-6}	18	30	180	307

^aSolar metallicity is assumed




FIG. 1.— Configuration of our CI Aql model in the late phases of the 2000 outburst. The cool component (*right*) is a slightly evolved MS companion ($1.5M_{\odot}$) filling up its inner critical Roche lobe. The north and south polar areas of the cool component are irradiated by the hot component ($1.2 M_{\odot}$ white dwarf, *left*). Here, we assume a non-axisymmetric accretion disk in order to reproduce asymmetric light curves as shown in Figure 3. The separation is $a = 4.25R_{\odot}$; the effective radii of the inner critical Roche lobes are $R_1^* = 1.53R_{\odot}$, and $R_2^* = R_2 = 1.69R_{\odot}$, for the primary white dwarf and the secondary main-sequence companion, respectively. The radius of the white dwarf in the cooling phase is as small as $R_{WD} = 0.0068R_{\odot}$, about 1/200 times the Roche lobe size. The WD radius is exaggerated in this figure so that it may be seen easily.




FIG. 2.— Calculated V light curves are plotted against the binary phase (binary phase is repeated twice from -1.0 to 1.0) together with the observational points of the two late phases (filled circles are in April/May of 2001, and small open circles are in August of 2001), and the quiescent phase (large open circles: smoothly averaged by us from Mennickent & Honeycutt (1995)) as well as the revised quiescent phase data (middle size open circles: Honeycutt 2002, private communication). The visual magnitude in quiescence has been revised by 0.62 mag down as seen in the figure. Here we adopted the ephemeris given by Matsumoto et al. (2001) as in equation (11). The model is a binary system of $1.2M_{\odot}$ white dwarf (WD) + $1.0M_{\odot}$ main-sequence (MS) companion. The other model parameters are shown in the figure and Table 1. The solid line for August of 2001 indicates the case of asymmetric structure of accretion disk while the dashed line denotes the case of symmetric configuration of accretion disk. The other solid lines correspond to the symmetric accretion disks. Here “asymmetry=0.75” means the value of $\zeta_{\text{low}} = 0.75$.




FIG. 3.— Same as those in Fig.2, but for a binary system of $1.2M_{\odot}$ WD + $1.5M_{\odot}$ MS companion.




FIG. 4.— Same as those in Fig.2, but for a binary system of $1.2 M_{\odot}$ WD + $2.0 M_{\odot}$ MS companion.




FIG. 5.— Same as those in Fig.3, but for an enlargement around eclipse minima. Only the data of August 2001 are shown together with the model light curves, solid line for asymmetric and dashed one for symmetric configurations of the accretion disk.




FIG. 6.— Same as those in Fig.1, but for the configuration at the eclipse minimum.




FIG. 7.— Calculated $B - V$ light curve in quiescence is plotted against the orbital phase for a pair of $1.2 M_{\odot}$ WD and $1.5 M_{\odot}$ MS companion.




FIG. 8.— Calculated B light curves are plotted against time (HJD 2,421,000+) together with the observational points of the 1917 outburst. The WD mass is attached to each light curve. Open circles indicate observational points (taken from Williams 2000). The model consists of a bloated white dwarf photosphere with no accretion disk and a nonirradiated main-sequence companion. The hydrogen content of the white dwarf envelope is assumed to be $X = 0.70$ for all models. The apparent distance modulus of $(m - M)_B = 15.06$ is assumed for all the white dwarf masses. Solid lines indicate the light curves connecting the B light at the binary phase 0.5 (roughly the mean brightness) while dashed lines correspond to those connecting the B light at the binary phase 0.0 (mid eclipse).




FIG. 9.— Calculated V , B , and I_c light curves are plotted against time (HJD 2,450,000+) together with the observational points of the CI Aql 2000 outburst. Small dots (small filled circles, small open circles) indicate observational V and visual magnitudes, while open squares represent observational B magnitudes and large open circles indicate observational I_c magnitudes (all taken from VSNET archives). Two V -magnitudes in the late-phase observed by Schaefer (2001a,b) are indicated by $\odot$ marks. Each light curve connects the brightness at the binary phase 0.4. The apparent distance moduli are $(m-M)_V = 14.06$, $(m-M)_B = 15.06$ and $(m-M)_I = 12.46$ for the V , B , and I_c light curves, respectively.